

349

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
18 November 2004 (18.11.2004)

PCT

(10) International Publication Number
WO 2004/098461 A2

(51) International Patent Classification: **A61F 2/04**

(21) International Application Number:
PCT/US2004/013502

(22) International Filing Date: 30 April 2004 (30.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
RM2003A000210 30 April 2003 (30.04.2003) IT

(71) Applicant (for all designated States except US): **AMS RE-
SEARCH CORPORATION** [US/US]; 10700 Bren Road
West, Minnetonka, MN 55343 (US).

(72) Inventor: **CERVIGNI, Mauro**; 10700 Bren Road West,
Minnetonka, MN 55343 (US).

(74) Agent: **HOHENSHELL, Jeffrey, J.**; 10700 Bren Road
West, Minnetonka, MN 55343 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).


Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: PROSTHESIS FOR USE IN THE SURGICAL TREATMENT OF UROGENITAL PROLAPSE AND FEMALE URI-
NARY INCONTINENCE

(57) Abstract: A surgical prosthesis is disclosed.



WO 2004/098461 A2

**PROSTHESIS FOR USE IN THE SURGICAL TREATMENT OF UROGENITAL
PROLAPSE AND FEMALE URINARY INCONTINENCE**

[0001] Cross-Reference to Related Applications

[0002] The present application claims the benefit of Italian Patent Application No. RM2003A000210 filed April 30, 2003, entitled, "Protesi da impiegarsi nella terapia chirurgica del prollasso urogenitale e dell'incontinenza urinaria femminile", inventor Mauro Cervigni (the entire contents incorporated by reference herein).

[0003] Background

[0004] The invention concerns a prosthesis 1 which consists of a mesh made of non-absorbable or partially absorbable or biologic material. The central body 2 has a first portion 3 designed to be placed under the bladder neck B and the middle urethra C, and a second portion 4 to be applied to the body D and the base E of the bladder beneath the inferior part of the bladder; the portions are nominated first and second as this is the sequence of the direction of insertion of the prosthesis; a first pair of arms 5 and 6 which extend from each side of the first portion 3 of the central body 2, the said arms 5 and 6 designed to be placed in the obturator foramen F bilaterally and a second pair of arms 7 and 8 which extend from each side of the second section 4 of the central body 2, the said arms 7 and 8 designed to be applied to the lateral perineum to the labia major and laterally to the bulbocavernosus muscles.

[0005] SUMMARY OF THE INVENTION

[0006] The present invention concerns a prosthesis for use in the surgical treatment of urogenital prolapse and female urinary incontinence.

[0007] More particularly the invention refers to a prosthesis to be used concomitantly in the surgical treatment of urogenital prolapse and female urinary incontinence allowing both problems to be resolved with a single operation.

[0008] The term urogenital prolapse refers to, in general, with reference to the physiology of the female pelvis, the inferior displacement of the uterus, the bladder and the rectum with consequent dislocation of the vaginal walls.

[0009] During the course of biological evolution, the humans assumed an erect posture which has determined, from the anatomical point of view, reversal of the position of the viscera inside the so-called pelvic cavity.

[0010] In particular, the female pelvis must support the weight of the intraabdominal organs, without adequate counter-support from the pelvic structures.

[0011] In this context, the above-mentioned urogenital prolapse can be caused by many factors, amongst which any phenomenon which increases the intraabdominal pressure on the female pelvis, for example, traumatic sporting activity, long periods of time standing up, chronic cough, and obesity.

[0012] In addition, prolapse is more common in women who have had multiple deliveries but also where there is weakness and loss of tone in the structures which support the uterus, even in nulliparous women.

[0013] The consequences of a prolapse of lesser or greater severity lead to a general deterioration in the quality of life and provoke local circulatory disturbances, swelling of the external genitalia, back pain, feeling of weight in the lower abdomen and in serious cases ulceration and exudation and difficulty with micturition and defecation.

[0014] Moreover, an important consequence of prolapse is stress urinary incontinence, due to the descent of the anterior wall of the vagina, the urethra and the bladder neck.

[0015] Various types of urogenital prolapse may be distinguished, depending on the area involved, its precise position in this area and the stage of descent, that is: Stage I when the lower limit of the uterine cervix or the anterior vaginal wall reaches the middle

of the vaginal canal without reaching the vulval orifice; Stage II when the uterine cervix or the anterior vaginal wall reaches the vulval orifice; Stage III when the lower extremity of the uterine cervix or the anterior vaginal wall protrudes from the vulva.

[0016] The term hysterocele refers to prolapse of the uterus; urethrocele when the urethra is prolapsed; cystocele when the bladder descends; rectocele when the rectum descends. In cases where more than one condition is present the terms are combined, for example urethro-cystocele or recto-cystocele.

[0017] A cystocele can occur because of detachment of the lateral or transverse endopelvic fascia or both (centro-lateral).

[0018] These different modalities are treated with specific surgical approaches which at times differ amongst themselves.

[0019] Operations for repair of lateral detachment of the fascia have been proposed, both abdominal (paravaginal repair) and vaginal (vaginal paravaginal repair); four points of the anterior vaginal wall may be surgically attached with four pairs of non-absorbable sutures to the retropubic area, in the so-called "four corner" technique. In central detachments, the approach consists of medializing the adjacent vaginal structures, in other words, the levator muscles and fascia, to raise the vagina, the bladder and sometimes the uterus.

[0020] With regards to the rectocele, the surgical approach includes the so-called posterior colporrhaphy, in other words the anchorage of the involved tissues to the medial part of the levator ani (Levator Myorrhaphy), or transanal treatment.

[0021] In spite of this variety of surgical therapies which allow the urogenital prolapse to be treated even at an early stage, when the various modalities are compared, the results are not satisfactory as there is a high rate of recurrence, up to 30%, after a follow-up period of 6 months to one year after the operation.

[0022] A recent analysis of data from the international literature has demonstrated that there is a recurrence rate of 20 to 78% for surgical repair of anterior vaginal wall prolapse with a follow-up of 6-48 months.

menopause; repeated pelvic surgery, for example hysterectomy. From a pathophysiological point of view the female urethral sphincter which is already weak is further weakened by stretching of the ligaments of support and relaxation of the perineal muscles (anal levator muscles).

[0031] In the past the various techniques for correcting SUI used the patient's own tissue to reinforce the sphincteric activity. A careful analysis of the literature demonstrates, however, that the recurrence rate can reach 15-20%. Thanks to the work of the Swedish School (Ulmsten) a new mini-invasive surgical procedure using Prolene® mesh has been introduced, applied with a "tension-free" technique under the middle-third of the urethra. The two ends of this tape are then transferred to the suprapubic area by 2 curved needles. A more recent variant uses the obturator foramen to stabilize the mesh under the urethra. In this way the support of the urethral sphincter is facilitated as "physiologically" as possible.

[0032] At the present time therefore a prosthesis is available for the treatment of SUI and a prosthesis is available for the treatment of genital prolapse but separate operations are required to resolve the above problems with consequent inconvenience for the patient. It is thus evident that there is a need for a new prosthesis which could resolve concurrently the problem of genital prolapse and SUI in a single operation.

[0033] The author of the present invention has prepared a prosthesis consisting of a mesh made of non-absorbable or partially absorbable material to apply using a "tension-free" technique, i.e., without forces of traction on the surrounding tissues, able to resolve both the above mentioned problems.

[0034] Thus this invention is a prosthesis to be used in the surgical treatment of urogenital prolapse and female urinary incontinence, constituting a mesh made of non-absorbable or partially absorbable or biological material. The central body has a first portion designed to be placed adjacent to bladder under the bladder neck and middle urethra; a second portion is to be placed in relation to the body and base of the bladder, under the inferior part of the bladder. The portions are described as first and second as this is the sequence of the direction of insertion of the prosthesis; a first pair of arms which extend from each side of the first portion of the central body, the said arms designed to be placed in the obturator foramen bilaterally and a second pair of arms

[0023] It is just as frequent that a patient treated for a prolapse in one area develops over time a prolapse in the non-treated area.

[0024] The cause of recurrence may be principally explained by the fact that any operation for repair demands, in the period of scar formation, a stabilization and reinforcement of the relevant connective structures. In reality, in many of these cases, the newly formed collagen constitutes a predisposition to successive descent of the surgically treated structures.

[0025] In fact, the connective tissue is weakened especially because in the typical age in which prolapse occurs (peri- or post-menopausal women) the fall in estrogens favors the formation of a weaker type of collagenous tissue.

[0026] From 1995 the Author of the present Invention has introduced the use of synthetic mesh in the repair of genital prolapse, taking inspiration from the procedure proposed and carried out in General Surgery, that is to say the use of synthetic mesh in the repair of inguinal hernias. The recurrence rate of inguinal hernias has consequently fallen from 20-30% to 1-2%.

[0027] Equally the use of the synthetic mesh for repair of genital prolapse has drastically reduced the recurrence rate from 30-50% in the past to the present 3-6%.

[0028] The materials used are varied and the "ideal" material has not yet been identified, but Prolene® (interwoven synthetic monofilament: Polypropylene) comes closest to the ideal requisites of a biocompatible material. These materials, in order to reduce complications and long term sequelae to a minimum, must be applied "tension free", in the sense that they must not cause traction or tension on the vaginal wall or bladder or rectum with consequent inevitable alteration of the functioning of these organs.

[0029] Often urogenital prolapse is also associated with urinary incontinence.

[0030] Stress urinary incontinence (SUI) is a condition that predominately affects women and is characterized by uncontrolled loss of urine during physical activity, or stress such as lifting a heavy object. The causes are various, amongst which: labor and delivery which determine trauma during the passage of the fetus in the vaginal canal;

which extend from each side of the second section of the central body, the said arms designed to be placed on the perineum, laterally to the labia major and passing through the bulbocavernosus muscles.

[0035] Preferably, the above named arms designated as first and second pair, are intended, at the respected ends not joined to the central body, to be a means for the attachment of single-use curved needles, that is the so-called means of attachment can be non-reusable in nature.

[0036] In particular, the said arms of the first pair of arms are angled with respect to the central body of the prosthesis in the opposite direction to the said direction of insertion of the actual prosthesis.

[0037] Continuing the description of the invention, the said arms of the second pair of arms are angled with respect to the central body of the prosthesis in the aforesaid direction of insertion of the prosthesis. Preferably, both the first and second pair of arms are coated with a nylon sheath to assist sliding over the tissues.

[0038] The prosthesis according to the present invention is made of a synthetic non-absorbable or partially absorbable. The present invention concerns a prosthesis for use in the surgical treatment of urogenital prolapse and female urinary incontinence.

[0039] The mesh, sterile and inert, which has bidirectional elastic extensibility and a resistance to traction greater than 10 kg/cm², being from 0.5 mm to 0.9 mm thick, preferably about 0.7 mm.

[0040] Regarding the intersections between the threads of the mesh, these are interconnected so that the mesh can be cut in any direction without unraveling.

[0041] The transverse dimensions of the pores, approximately rhomboid shaped, are between 1 mm and 3 mm.

[0042] Each thread is a polypropylene (C₃H₆) non-absorbable monofilament, more specifically an isotactic stereoisomer of polypropylene.

[0043] A type of mesh, which is particularly suitable for this type of application, is a mesh made of Prolene® monofilament, made by Ethicon® Inc. and marketed under the commercial name of polypropylene PROLENE® mesh.

[0044] As an alternative to the non-absorbable mesh described above, it is possible to use with the same efficacy a partially absorbable mesh or a biological material such as dermis, pericardium, or intestinal submucosa. Preferably, the mesh should be made of biological tissue in the central part and synthetic material in both the first and second pair of arms.

[0045] The surgical technique is now described below.

[0046] A longitudinal incision is made from 1 cm below the external urethral meatus to the anterior vaginal fornix. The underlying pubocervical fascia is then detached, arriving laterally to the ascending branch of the ischiopubic and laterally to the lateral fornix. Two (2) incisions are then carried out 1 cm above the clitoris in the inguinal folds and a curved needle is passed through the obturator foramen and exits at the sides of the bladder neck bilaterally. The apex of an arm of the prosthesis is then threaded into the needle (or attached using a Luer-Lock) and then pulled through the previous track. Two (2) further incisions of 1-2 cm are then executed laterally to the labia major at the level of the vaginal vestibule and with the same curved needle the bulbo-cavernosous muscle is penetrated and the end of the needle is brought out 2 cm posterior to the lateral fornix. A free flap of the mesh is placed under the base of the bladder and the vaginal opening is closed with a continuous suture. With a combined maneuver, applying traction to the arms which have been placed outside and on the anterior vaginal wall, the mesh is positioned as gently as possible, the excess tissue is then excised and the four (4) cutaneous openings are closed.

[0047] New findings regarding the etiology of pelvic descent have meant that the cystocele is considered to be a true "bladder hernia". The alteration of and relaxation of the endopelvic fascia ("The anatomy of the pelvic floor", J. O. DeLancey, Curr. Opin. Obstet. Gynecol. 1994 Aug; 6(4): 313-6) and the sagging of the muscles and ligaments ("Anatomy and biomechanics of genital prolapse", J. O. DeLancey, Clin. Obstet. Gynecol. 1993 Dec; 36(4): 897-909) are the cause of the initial process of "bladder herniation". The therapeutic strategy thus tends to be analogous to that of the hernia.

The positioning of a "tension free" mesh tends to restore the altered support of the endopelvic fascia. The concept of the "tension free" technique has the further advantage of respecting the lines of force acting at the pelvic exit thus avoiding anatomical distortion, rather frequent in the techniques of repair in pelvic surgery. For these reasons the use of "tension free" meshes placed transperineally can be considered as a new technique in the correction of cystocele as well as Stress Urinary Incontinence (SUI).

[0048] The prosthesis according to the present invention has been positioned by the author using the technique described above in a large number of patients, about 400 with a follow-up of greater than 3 years, obtaining a cure rate of more than 90% with a low incidence of complications including: erosion (6%), dyspareunia (5%), pelvic pain (7%).

[0049] A further advantage of the prosthesis according to the present invention is an increase in the production of connective tissue between the pores of the surgically implanted mesh, creating a natural non-absorbable and elastic support which provides an effective support and prevents a further descent of the pelvic structures subject to prolapse and incontinence.

[0050] Moreover, the surgical application of the aforesaid prosthesis, which can be carried out both at an early and late stage of prolapse and incontinence, is able to correct all the existing defects, centrally and/or laterally and SUI and to prevent the potential appearance of further defects, so that a differential diagnosis of the position of the prolapse preparatory to surgical therapy is actually superfluous.

[0051] The present invention will now be described in a explanatory but not limitative document, according to the preferred modes of implementation, with particular reference to the diagrams attached, in which:

[0052] Figure 1 shows a form of the prosthesis according to the present invention;

[0053] Figure 2 shows a schematic design of the prosthesis of figure 1 applied to the female urogenital apparatus seen posteriorly;

[0054] Figure 3 shows a schematic design of the prosthesis of figure 1 applied to the female urogenital apparatus seen anteriorly;

[0055] Figure 4 shows the final phase of the surgical technique for the application of the prosthesis of figure 1 in which the incisions for insertion of the arms of the prosthesis are illustrated.

[0056] Observing the figures of the attached diagrams, a form of realization of the prosthesis according to the invention is demonstrated, indicated generically with the reference number 1.

[0057] The prosthesis 1 constitutes a mesh made of non-absorbable or partially absorbable or biological material and includes a central body 2 with a first portion 3 designed to be placed under the bladder neck B and the middle urethra C, and a second portion 4 to be applied to the body D and the base E of the bladder beneath the inferior part of the bladder; the portions are nominated first and second as this is a preferred sequence of the direction of insertion of the prosthesis; a first pair of arms 5 and 6 which extend from each side of the first portion 3 of the central body 2, the said arms 5 and 6 designed to be placed in the obturator foramen F bilaterally and a second pair of arms 7 and 8 which extend from each side of the second section 4 of the central body 2, the said arms 7 and 8 designed to be applied to the perineum lateral to the labia major and penetrating the bulbocavernosus muscles (see figure 4 which demonstrates the incisions H through which pass arms 7 and 8). To facilitate the introduction and fixation of the 4 arms, the latter are covered with a nylon sheath which is then withdrawn once placed in position.

[0058] As described above, the surgical technique for the application of the prosthesis according with the present invention essentially consists of exposing the anterior vaginal wall and making a longitudinal incision from 1 cm below the external urethral meatus to the anterior vaginal fornix. The underlying pubocervical fascia is then detached, as far as the ascending branch of the ischiopubic and laterally to the lateral fornix. Two (2) lateral incisions (G) are then carried out 1 cm above the clitoris in the inguinal folds and a curved needle is passed through the obturator foramen (F) and exits at the sides of the bladder neck bilaterally. The apex of an arm of the prosthesis is then threaded into the needle and then pulled through the previous track, in this way positioning the first portion (3) of the central body (2) next to the bladder (A) under the bladder neck (B) and middle urethra (C) and the arms (5 and 6) in the obturator foramen

(F) (see figure 4 which shows the incisions G through which arms 5 and 6 pass). Two (2) further incisions (H) of 1-2 cm are then executed laterally to the labia major at the level of the vaginal vestibule and with the same curved needle the track is penetrated, laterally to the bulbo-cavernosous muscle and the end of the needle is brought out 2 cm posterior to the lateral fornix, in order to position the second portion (4) of the central body (2) of the prosthesis (1) next to the bladder body (D) and the bladder base (E) and under the inferior part of the bladder and the arms (7 and 8) in the perineum laterally to the labia major and penetrating the perineum laterally to the bulbo-cavernosous muscle (see figure 4 which shows the incisions H through which arms 7 and 8 pass). A free flap of the mesh is placed under the inferior part of the bladder, and after having verified the lack of tension of the two anterior arms under the bladder neck using Metzenbaum scissors, the vaginal opening is closed with a continuous suture. With a combined maneuver, applying traction to the arms which have been placed outside and on the anterior vaginal wall (using two fingers placed in vagina), the mesh is positioned as gently as possible, the excess tissue is then excised and the 4 cutaneous openings are closed (G, H).

[0059] The present invention has been described in an explanatory but not limitative document, according to the preferred modes of realization, but it should be understood that variations and/or modifications may be added, without exceeding the relative protection, as defined by the attached claims.

CLAIMS

1. Prosthesis to be used in the surgical treatment of urogenital prolapse and female urinary incontinence constituting a mesh made of non-absorbable or partially absorbable or biological material, the prosthesis comprising a central body having a first portion designed to be placed beneath the bladder under the bladder neck and middle urethra; a second portion is to be placed in relation to the bladder, under the inferior part of the bladder, a first pair of arms which extend from each side of the first portion of the central body, the said arms designed to be placed in the obturator foramen bilaterally and a second pair of arms which extend from each side of the second portion of the central body, the said second arms designed to be placed on the perineum, laterally to the labia major and passing through the levator muscles.
2. Prosthesis according to claim 1 characterized in that the said arms of the said first and second pair are intended, at the respective ends not attached to the said central body, to be the means of attachment of curved surgical needles, preferably single use curved surgical needles.
3. Prosthesis according to claim 2 characterized in that the said means of attachment are non-reusable.
4. Prosthesis according to claim 2 characterized in that the said arms of the said first and second pair are covered with a nylon sheath.

5. Prosthesis according to claim 1 characterized in that the said arms of the first pair of arms are angled with respect to the central body of the prosthesis in the opposite direction to the preferred direction of insertion of this prosthesis.
6. Prosthesis according to claim 1 characterized in that the said arms of the second pair of arms are angled with respect to the central body of the prosthesis in the preferred direction of the insertion of this prosthesis.
7. Prosthesis according to claim 1 characterized in that the non-absorbable or partially absorbable synthetic mesh has bi-directional elastic distensibility and a resistance to traction greater than 10 KG/cm².
8. Prosthesis according to claim 7 characterized in that the mesh is between 0.5 mm and 0.9 mm thick.
9. Prosthesis according to claim 7 characterized in that the mesh is approximately 0.7 mm thick.
10. Prosthesis according claim 1 characterized in that the mesh is made of polypropylene.
11. Prosthesis according to claim 1 characterized in that the mesh is made of biological material centrally and synthetic material in the arms of both the first and second pairs of arms.

12. Prosthesis according to claim 2 characterized in that the said arms of the first pair of arms are angled with respect to the central body of the prosthesis in the opposite direction to the preferred direction of insertion of this prosthesis.

13. Prosthesis according to claim 3 characterized in that the said arms of the first pair of arms are angled with respect to the central body of the prosthesis in the opposite direction to the preferred direction of insertion of this prosthesis.

14. Prosthesis according to claim 4 characterized in that the said arms of the first pair of arms are angled with respect to the central body of the prosthesis in the opposite direction to the preferred direction of insertion of this prosthesis.

1/2




FIG.1




FIG.2

2/2




FIG. 3




FIG. 4

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
18 November 2004 (18.11.2004)

PCT

(10) International Publication Number
WO 2004/098461 A3

(51) International Patent Classification?: **A61F 2/00**

(21) International Application Number:
PCT/US2004/013502

(22) International Filing Date: 30 April 2004 (30.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
RM2003A000210 30 April 2003 (30.04.2003) IT

(71) Applicant (for all designated States except US): **AMS RE-
SEARCH CORPORATION** [US/US]; 10700 Bren Road
West, Minnetonka, MN 55343 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **CERVIGNI, Mauro**;
10700 Bren Road West, Minnetonka, MN 55343 (US).

(74) Agent: **HOHENSHELL, Jeffrey, J.**; 10700 Bren Road
West, Minnetonka, MN 55343 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,

CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).


Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(88) Date of publication of the international search report:
30 June 2005

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **PROSTHESIS FOR THE TREATMENT OF UROGENITAL PROLAPSE AND FEMALE URINARY INCONTI-
NENCE**



(57) Abstract: The invention concerns a prosthesis (1) which consists of a mesh made of non- absorbable or partially absorbable or biologic material. The central body has a first portion (3) designed to be placed under the bladder neck (B) and the middle urethra, and a second portion (4) to be applied to the body (D) and the base (E) of the bladder (A) beneath the inferior part of the bladder (A); the portions are nominated first and second as this is the sequence of the direction of insertion of the prosthesis (1); a first pair of arms (5,6) which extend from each side of the first portion (3) of the central body, the said arms (5,6) designed to be placed in the obturator foramen (F) bilaterally and a second pair of arms (7,8) which extend from each side of the second section (4) of the central body, the said arms (7,8) designed to be applied to the lateral perineum to the labia major and laterally to the bulbocavernosus muscles.

WO 2004/098461 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2004/013502

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61F2/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 774 240 A (ETHICON INC) 21 May 1997 (1997-05-21)	1-3, 5, 6, 8-14
Y	column 1, line 33 - column 5, line 36 claims; figure 1	4
A	WO 01/06951 A (ANGIOLOGICA B M S R L) 1 February 2001 (2001-02-01)	1-3, 5-14
Y	pages 3-4 claims 1-3, 7, 8; figures	4
A	US 6 042 534 A (GELLMAN BARRY N ET AL) 28 March 2000 (2000-03-28)	1
	claims; figures 10a-11	
A	US 2002/107430 A1 (NEISZ JOHANN J ET AL) 8 August 2002 (2002-08-08)	1-14
	claims; figures	
	----- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

9 May 2005

Date of mailing of the international search report

19/05/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3015

Authorized officer

Kuehne, H-C

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US2004/013502

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 02/38079 A (ETHICON GMBH) 16 May 2002 (2002-05-16) claims; figures -----	1-14
A	WO 00/27304 A (SOFRADIM PRODUCTION SA) 18 May 2000 (2000-05-18) claims; figures -----	1-14

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2004/013502

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0774240	A	21-05-1997	DE 19544162 C1	24-04-1997
			BR 9605581 A	18-08-1998
			CA 2190449 A1	18-05-1997
			DE 69626604 D1	17-04-2003
			DE 69626604 T2	27-11-2003
			EP 0774240 A1	21-05-1997
			JP 9201376 A	05-08-1997
			US 5840011 A	24-11-1998
WO 0106951	A	01-02-2001	IT MI991660 A1	29-01-2001
			AU 5987000 A	13-02-2001
			WO 0106951 A1	01-02-2001
US 6042534	A	28-03-2000	AU 6329598 A	08-09-1998
			CA 2280814 A1	20-08-1998
			DE 69808614 D1	14-11-2002
			DE 69808614 T2	18-06-2003
			EP 0983033 A1	08-03-2000
			JP 2001511685 T	14-08-2001
			WO 9835632 A1	20-08-1998
US 2002107430	A1	08-08-2002	AT 274329 T	15-09-2004
			CA 2403687 A1	01-08-2002
			CA 2434461 A1	01-08-2002
			CA 2434463 A1	01-08-2002
			DE 60105200 D1	30-09-2004
			EP 1353597 A1	22-10-2003
			EP 1353598 A1	22-10-2003
			EP 1353599 A2	22-10-2003
			EP 1508305 A2	23-02-2005
			US 2002147382 A1	10-10-2002
			WO 02058562 A1	01-08-2002
			WO 02058563 A1	01-08-2002
			WO 02058564 A2	01-08-2002
			US 2003050530 A1	13-03-2003
			US 2004015048 A1	22-01-2004
			US 2004015057 A1	22-01-2004
			US 2004068159 A1	08-04-2004
			US 2002151762 A1	17-10-2002
			BR 0116812 A	27-01-2004
			CA 2404459 A1	15-08-2002
			CA 2434027 A1	01-08-2002
			EP 1353600 A2	22-10-2003
			EP 1353601 A1	22-10-2003
			JP 2004526483 T	02-09-2004
			WO 02062237 A1	15-08-2002
			WO 02058565 A2	01-08-2002
			US 2003045774 A1	06-03-2003
			US 2003130670 A1	10-07-2003
			US 2004106845 A1	03-06-2004
			US 2002099258 A1	25-07-2002
			US 2002099259 A1	25-07-2002
WO 0238079	A	16-05-2002	DE 10056169 A1	06-06-2002
			AT 268146 T	15-06-2004
			AU 2952502 A	21-05-2002
			CA 2428652 A1	16-05-2002
			CN 1486164 A	31-03-2004

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2004/013502

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0238079	A		DE 60103660 D1	08-07-2004
			DE 60103660 T2	30-09-2004
			DK 1333776 T3	27-09-2004
			WO 0238079 A2	16-05-2002
			EP 1333776 A2	13-08-2003
			ES 2222403 T3	01-02-2005
			JP 2004522470 T	29-07-2004
			US 2004116774 A1	17-06-2004

WO 0027304	A	18-05-2000	FR 2785521 A1	12-05-2000
			EP 1128781 A1	05-09-2001
			WO 0027304 A1	18-05-2000
			US 6575897 B1	10-06-2003
